Analytical and Monte Carlo Studies of Jets with Heavy Mesons and Quarkonia

Reggie Bain
Duke University

In collaboration with L. Dai, A. Hornig, A.Leibovich, Y. Makris, T. Mehen

Quarkonium 2016, PNNL, Richland, WA
June 6-10, 2016

Motivations

Understand high energy jets at LHC

- *Testing QCD*
- *Calculating backgrounds for new physics*



ATLAS Collaboration

Study wealth of jet substructure observables



Thaler, v. Tilberg, arXiv:1011.2268

Elucidate outstanding puzzles in quarkonia production



LHCb Collaboration

Outline

- Fragmenting jet functions w/ angularities
- NLL' cross section calculations
  - $e^+e^- \longrightarrow$ *2 jets with B meson*
  - $e^+e^- \longrightarrow$ *3 jets with J/ψ from gluon*
- Comparisons of NLL' vs. Monte Carlo

Studying Jets w/ SCET

Factorization ⟷ Short Distance x Long Distance



Figure from "Lectures on the Soft Collinear Effective Theory" by Iain W. Stewart, 2013

Factorization Theorem

$$d\sigma = H \times J^1 \otimes J^2 \otimes S$$

- Hard function $H(\mu)$
- Jet Functions $J^{(1)}(\mu)$
- Soft Function $S(\mu)$

→ Measured

→ Unmeasured

Fragmenting Jet Functions (FJF's)

Measured Jets w/ Identified Hadron



Jet Inv. Mass	s
Plus Momentum	$p_H^+ = z p_J^+$
Cone Size	R

Closely related to Fragmentation Functions (FF's)

Probe jet invariant mass

$$D_i^h(z,\mu) \longrightarrow \mathcal{G}_i^h(s,z,\mu)$$

Procura, Stewart, arXiv:0911.4980

Generalize FJF Formalism

Measured Angularity of Jet

$$\tau_a = \frac{1}{\omega} \sum_i (p_i^+)^{1-a/2} (p_i^-)^{a/2}$$

Sum over jet particles i

$$\omega = \sum_i p_i^- \approx 2E_{jet}$$

a=0 ***jet invariant mass*** $s = \omega^2 \tau_0$

a=1 ***jet broadening***



We have good analytic handle on τ_a

Hornig, Lee, Ovanesyan, arXiv:0905.0168
S.D.Ellis, et. al (2010), arXiv:1001.0014

Cross-Section with FJFs

Jet cross-section ⟶ Jet w/ Identified Hadron cross-section

$$J_i(\tau_a, \mu) \to \frac{1}{2(2\pi)^3} \mathcal{G}_i^H(\tau_a, z, \mu) dz$$

Convolution of Matching Coefficients & Fragmentation Functions (FF's)

$$\mathcal{G}_i^H(\tau_a, z, \mu) = \sum_j \left[\mathcal{J}_{ij}(\tau_a, \mu) \bullet D_j^H(\mu) \right](z)$$

$$[f \bullet g](z) \equiv \int_z^1 \frac{dx}{x} f(x) g(z/x)$$

Calculate $\mathcal{J}_{ij}(\tau_a, z, \mu)$ perturbatively for different observables

Procura, Stewart, arXiv:0911.4980

Cross-sections for $e^+e^- \rightarrow$ jets

R. Bain, L. Dai, A. Hornig, A.Leibovich, Y. Makris, T. Mehen, arXiv:1603.06981

$e^+e^- \rightarrow b\bar{b}$ vs. Monte Carlo

↳ B jet

$e^+e^- \rightarrow b\,\bar{b}\,g$ vs. Monte Carlo

↳ J/ψ jet

Goals

1. Study z, τ_0 distributions
2. Compare NLL' with PYTHIA & HERWIG

Cross Section for 2 jets & B^+/B^0

Re-summed to NLL' using renormalization group (RG)

FF for B's at jet scale (DGLAP)

$$d\sigma(\tau_a, z) \equiv \frac{1}{\sigma_0}\frac{d\sigma^{(b)}}{d\tau_a dz} = H_2(\mu_H) \times S^{\text{unmeas}}(\mu_\Lambda) \times J_{\bar{n}}^{(\bar{b})}(\mu_{J_{\bar{n}}}) \times$$

$$\times \sum_j \left\{ \left(\frac{\Theta(\tau_a)}{\tau_a^{1+\Omega}}\right) \left[\delta_{bj}\delta(1-z)\left(1 + f_S(\tau_a, \mu_{S^{\text{meas}}})\right) + f_{\mathcal{J}}^{bj}(\tau_a, z, \mu_{J_n})\right] \bullet \frac{D_{j\to B}(z, \mu_{J_n})}{2(2\pi)^3} \right\}_+$$

$$\times \Pi(\mu, \mu_H, \mu_\Lambda, \mu_{J_{\bar{n}}}, \mu_{J_n}, \mu_{S^{\text{meas}}})$$

coupled z & τ_a

RG evolution factor

Coupling of z and τ_a dependence appears first at NLO

Evaluate each piece at characteristic scale, evolve up to hard scale

b quark Fragmentation Function

Fit power model to LEP data

Inclusive Cross-Section vs. z



Fragmentation Function

$$D(x, \mu_0) = N x^{\alpha}(1-x)^{\beta}$$

$$N = 4684.1$$

$$\alpha = 16.87$$

$$\beta = 2.028$$

$$\mu_0 = m_b = 4.5 GeV$$

$$\chi^2_{dof} = 1.495$$

Kniehl, et al. arXiv:0705.4392

NLL' vs. Monte Carlo (B^+/B^0)

z distributions for fixed τ_a match well



Monte Carlo Details

SISCone, $R = 0.6$, $N_{jets} = 2$

$$E_{jet} > \frac{E_{cm} - \Lambda}{N_{jets}}, \quad E_{cm} = 250\text{GeV}$$

$\Lambda = 30\text{GeV}$

NLL' vs. Monte Carlo (B^+/B^0)

τ_0 distributions for fixed z also show good agreement







Monte Carlo Details

SISCone, $R = 0.6$, $N_{jets} = 2$

$$E_{jet} > \frac{E_{cm} - \Lambda}{N_{jets}}, \quad E_{cm} = 250\text{GeV}$$

$\Lambda = 30\text{GeV}$

Next steps

- NLL' and Monte Carlo match well for B mesons in jets

From B's to J/ψ's…

- FJF's discriminate between J/ψ production mechanisms

Baumgart, Mehen, Leibovich, Rothstein, arXiv: 1406.2295 (2014)

- Calculate 3 jet cross-section with J/ψ, compare with MC…

NRQCD Fragmentation Functions

Fragmentation Function vs. z of J/ψ



NRQCD Factorization

$$D_{g\to J/\psi} = \sum_n D^{(n)}_{g\to J/\psi} \langle \mathcal{O}^{J/\psi}(n)\rangle$$

with $n = {}^{2S+1}L_J^{(1,8)}$

Examples of $\alpha_s(2m_c)$ & z dependence

$$D^{^3S_1^{(8)}}_{g\to J/\psi}(z, 2m_c) = \frac{\pi\alpha_s(2m_c)}{24m_c^3}\delta(1-z)$$

$$D^{^1S_0^{(8)}}_{g\to J/\psi}(z, 2m_c) = \frac{5\alpha_s(2m_c)}{96m_c^3}\left(3z - 2z^2 + 2(1-z)\log(1-z)\right)$$

Braaten, Chen, hep-ph/9610401
Braaten, Chen, hep-ph/9604237
Braaten, Yuan, hep-ph/9302307

Extract LDME's from World's Data

Need CS+CO at NLO to fit data from various experiments





Fit to world data (2/26 plots shown) to $e^+e^-,\ \gamma\gamma,\ \gamma p,\ p\bar{p}, pp \to J/\psi + X$

$\langle \mathcal{O}^{J/\psi}(^3S_1^{(1)})\rangle$	$\langle \mathcal{O}^{J/\psi}(^3S_1^{(8)})\rangle$	$\langle \mathcal{O}^{J/\psi}(^1S_0^{(8)})\rangle$	$\langle \mathcal{O}^{J/\psi}(^3P_J^{(8)})\rangle/m_c^2$
1.32 GeV^3	2.24 $\times 10^{-3}$ GeV^3	4.97×10^{-2} GeV^3	-7.16 $\times 10^{-3}$ GeV^3

Buttenschon, Kniehl (2011), arXiv:1105.0820

NLL' vs. PYTHIA (J/ψ)

Monte Carlo/NLL' τ_a distributions for fixed z's show similarities



As z ⟶ 0 we see less dependence on production mechanism

Comparing NLL' & PYTHIA

PYTHIA yields much harder z distributions



Gluon Fragmentation and PYTHIA

PYTHIA's picture of showering off onia different from theory

Analytic



PYTHIA



Monte carlo z distributions much harder than analytic

Gluon Fragmentation Improved PYTHIA (GFIP)

Madgraph 5

$e^+e^- \to b\bar{b}g$



PYTHIA + Convolution



2. PYTHIA ⟶ No hadronization, adjust shower pT cutoff

3. Convolve NRQCD FFs w/ random final state gluon

Comparing NLL', PYTHIA, and GFIP

GFIP shows far better agreement w/ NLL'



Conclusions

- New calculation: FJF for measured angularities
- Our calculation fits B production in Monte Carlo ($d\sigma/d\tau dz$)
- Default Monte Carlo J/ψ seems to lack proper onia showering
- GFIP shows improvements in z-distributions

Current & Future Work

- In talks with PYTHIA experts on modifications to parton shower
- Calculate cross-section for pp w/ measured angularity
- Extend FJF formalism to other jet-substructure observables

Jet-shape

$$\Psi_J(r) \equiv \frac{\sum_{i,\,\theta_{i\hat{n}<r}} E^i}{\sum_{i,\,\theta_{i\hat{n}<R}} E^i} = \frac{E_r}{E_R}$$

pT and Δ_θ



Thank you!

Backup Slides

Minimizing Large Logs of 1-z

Use z-dependent measured jet scale in FJF ⟶ decreases uncertainty







Reorganize Logs of z

$\left[f_{\mathcal{J}}^{ij} \bullet D\right](z)$ contains $\log\left(\frac{\mu}{\mu_J(\tau, z)}\right)$

where

$$\mu_J(z) = \omega \tau^{1/(2-a)} (1-z)^{(1-a)/(2-a)}$$

Abbate, Fickinger, Hoang, Mateu, Stewart, arXiv:1006.3080
Hornig, Makris, Mehen, arXiv:1601.01319
Ligeti, Stewart, Tackmann, arXiv:0807.1926

Procura, Waalewijn, arXiv:1111.6605

Matching Coefficients at NLO

We calculated all 4 NLO (1-loop) $\mathcal{J}_{ij}$ for measured angularities

$$
\begin{aligned}
\frac{\mathcal{J}_{qq}(\omega, z, \tau_a, \mu)}{2(2\pi)^3} = & \frac{C_F \alpha_s}{2\pi} \frac{1}{\omega^2} \Bigg\{ \delta(\tau_a)\delta(1-z) \frac{2-a}{1-a} \left(-\frac{\pi^2}{12} + \frac{1}{2} \ln^2 \left(\frac{\mu^2}{\omega^2} \right) \right) \\
& + \delta(\tau_a) \left(1 - z - \left[\ln \left(\frac{\mu^2}{\omega^2} \right) + \frac{1}{1-a/2} \ln \left(1 + \frac{(1-z)^{1-a}}{z^{1-a}} \right) \right] \frac{1+z^2}{(1-z)_+} \right. \\
& \left. + \frac{1-a}{1-a/2} (1+z^2) \left(\frac{\ln(1-z)}{1-z} \right)_+ \right) \\
& + \left[\frac{1}{\tau_a} \right]_+ \left(\frac{1}{1-a/2} \frac{1+z^2}{(1-z)_+} - \delta(1-z) \frac{2}{1-a} \ln \left(\frac{\mu^2}{\omega^2} \right) \right) \\
& + \frac{2\delta(1-z)}{(1-a)(1-a/2)} \left[\frac{\ln \tau_a}{\tau_a} \right]_+ \Bigg\}
\end{aligned}
$$

also… $\mathcal{J}_{qg}, \mathcal{J}_{gq}, \mathcal{J}_{gg}$

Consistency checks

1. $\lim_{a \to 0} \mathcal{J}_{ij}(\tau_a, z, \mu) = \omega^2 \mathcal{J}_{ij}(s, z, \mu) \longrightarrow$ Jain, et. al, arXiv:1101.4953
2. $J_i(s, \mu) = \frac{1}{2(2\pi)^3} \sum_j \int_0^1 dz z \mathcal{J}_{ij}(s, z, \mu) \longrightarrow$ S.D.Ellis, et. al, arXiv:1001.0014

Cross Section for 3 jets & J/ψ

Re-summed to NLL' using renormalization group (RG)

NRQCD FF for J/ψ's

$$\frac{1}{\sigma_0}\frac{d\sigma^{(g)}}{d\tau_a dz} = H_3(\mu_H) \times S^{\text{unmeas}}(\mu_\Lambda) \times J_{n_1}^{(\bar{b})}(\mu_{J_{n_1}}) \times J_{n_2}^{(b)}(\mu_{J_{n_2}})$$

$$\times \sum_i \left\{ \left(\frac{\Theta(\tau_a)}{\tau_a^{1+\Omega}}\right) \left[\delta_{gi}\delta(1-z)(1+f_S(\tau_a, z, \mu_{S^{\text{meas}}})) + f_{\mathcal{J}}^{gi}(\tau_a, z, \mu_{J_{n_3}})\right] \bullet \frac{D_{i\to J/\psi}(z, \mu_{J_{n_3}})}{2(2\pi)^3} \right\}_+$$

$$\times \Pi(\mu, \mu_H, \mu_\Lambda, \mu_{J_{n_1}}, \mu_{J_{n_2}}, \mu_{J_{n_3}}, \mu_{S^{\text{meas}}}),$$

coupled z & τ_a

RG evolution factor

Similar to cross-section for B's with new FF and 3 jets

Want to study pp ⟶ focus on g fragmentation

NLL' vs. PYTHIA (J/ψ)

Monte Carlo/NLL' τ_a distributions for different a's also show similarities



More discriminating power for larger a (a < 1 in $SCET_I$)

Default MadGraph + PYTHIA

1. MadOnia: Create J/psi in hard process

$e^+e^- \to b\bar{b}\,c\bar{c}\left[{}^3S_1^{(8)}\right]$ $e^+e^- \to b\bar{b}\,g\,c\bar{c}\left[{}^1S_0^{(8)}\right]$ $e^+e^- \to b\bar{b}\,g\,g\,c\bar{c}\left[{}^3S_1^{(1)}\right]$



2. PYTHIA ⟶ Parton shower + hadronization

3. RIVET ⟶ Reconstruct jets + implement cuts

Comparing NLL', PYTHIA, and GFIP

GFIP shows far better agreement w/ NLL'



Extra Details on Scales/FJF's

Characteristic Scales in B meson case

Function (F)	H_2	$J^{\bar{b}}_{\bar{n}}$	S^{unmeas}	$\mathcal{J}(\tau, z)$	$S^{\text{meas}}(\tau)$
Scale (μ_F)	E_{cm}	$\omega_{\bar{n}} r$	$2\Lambda r^{1/2}$	$\omega_n \tau^{1/(2-a)}(1-z)^{(1-a)/(2-a)}$	$\omega_n \tau / r^{1-a}$

Previous Studies on FJF's

Different identified hadrons/measured observables

X. Liu, arXiv:1011.3872
Jain, Procura, Waalewijn, arXiv: 1101.4953
Jain, Procura, Waalewijn, arXiv:1110.0839
Procura, Waalewijn, arXiv:1111.6605
Jain, Procura, Waalewijn, B. Shotwell, arXiv:1207.4788
Bauer, Mereghetti, arXiv:1312.5605
Baumgart, Mehen, Leibovich, Rothstein, arXiv:1406.2295
Chien, Z.-B. Kang, F. Ringer, I. Vitev and H. Xing, arXiv:1512.06851

Profile Functions

Abbate, Fickinger, Hoang, Mateu, Stewart, arXiv:1006.3080
Ligeti, Stewart, Tackmann, arXiv:0807.1926
Hornig, Makris, Mehen, arXiv:1601.01319



$$\mu_S^{PF}(\tau) = \left[1 + \epsilon_S \frac{g(\tau)}{g(1)}\right] \times \begin{cases} \mu_0 + \alpha\tau^\beta; & 0 < \tau < \tau_{min} \\ \omega\tau/r^{(1-a)}; & \tau_{min} \leq \tau \end{cases}$$

$$\mu_J^{PF}(\tau) = \left[1 + \epsilon_J \frac{g(\tau)}{g(1)}\right] \times \begin{cases} (\omega r)^{(1-a)/(2-a)}(\mu_0 + \alpha\tau^\beta)^{1/(2-a)}; & 0 < \tau < \tau_{min} \\ \omega\tau^{1/(2-a)}; & \tau_{min} \leq \tau \end{cases},$$

Reorganizing Log(1-z)

Convolution in z

$$\frac{1}{T_{ij}}\frac{2\pi}{\alpha_s(\mu)} f^{ij}_{\mathcal{J}}(\tau,z,\mu) \bullet D(z) = \delta_{ij}\ f_1(\tau,z,\mu)\ D(z) - \int_z^1 dx\ \ f_2(\tau,x,\mu)\Big(\frac{\bar{P}_{ji}(x)}{x} \circ D\left(\frac{z}{x}\right)\Big)$$
$$+ \int_z^1 dx\ \ \Big[c_{ij}(x) - \frac{1}{1-a/2}\ln\left(1+\left(\frac{1-x}{x}\right)^{1-a}\right)\frac{\bar{P}_{ji}(x)}{x}\Big] \circ D\left(\frac{z}{x}\right)\,,$$

Definitions of functions

$$f_1(\tau,z,\mu) = \frac{1-a/2}{1-a}\Big(f_2(\tau,z,\mu)\Big)^2 + \frac{a(1-a/4)}{(1-a)(1-a/2)}\frac{\pi^2}{6} - \frac{1}{(1-a)(1-a/2)}\psi^{(1)}(-\Omega)$$

$$f_2(\tau,z,\mu) = 2\ln\left(\frac{\mu}{\mu_J(\tau,z)}\right) + \frac{1}{1-a/2}H(-1-\Omega)\,,$$

z dependent scale

$$\mu_J(\tau,z) = \omega\tau^{1/(2-a)}(1-z)^{(1-a)/(2-a)}$$

Splitting Kernels

* $$\bar{P}_{qq} = P_{qq} - \frac{3}{2}\delta(1-z) = \frac{1+z^2}{(1-z)_+}\,,$$

$$\bar{P}_{gq} = P_{gq} = \frac{1+(1-x)^2}{x}\,,$$

$$\bar{P}_{qg} = P_{qg} = x^2 + (1-x)^2\,,$$

* $$\bar{P}_{gg} = P_{gg} = 2\frac{(1-x+x^2)^2}{x(1-x)_+}\,,$$

$$\bar{P}_{ij}|_{i,j\neq q} = P_{ij}$$



* **Pythia evolution kernel**

* **Analytic evolution kernel**

Polarization Problem



λ_θ = +1 (trans.), 0 (unpol.), -1 (long.)

θ = J/ψ and μ+ momentum polar angle

Blue = No feed down, pT > 3 GeV; Buttenschon et. al (2012)

Red = Chi_cJ and Psi(2S) feed down, pT > 7 GeV; Gong et al. (2013)

Green = No feed down, pT > 7 GeV; Chao et. al (2012)

Magenta = Color singlet at NLO; Buttenschon et al (2012)

Terms that Arise at NLL'

Measured jet function contribution (NLO/NLL')

$$f_{\mathcal{J}}^{ij}(\tau, z, \mu) = T_{ij} \frac{\alpha_s(\mu)}{2\pi} \Bigg(c_0^{ij}(z, \mu) + c_1^{ij}(z, \mu) \Big(\ln \tau - H(-1-\Omega) \Big) + c_2 \delta_{ij} \delta(1-z) \Big(\frac{(\ln \tau - H(-1-\Omega))^2 + \pi^2/6 - \psi^{(1)}(-\Omega)}{2} \Big) \Bigg).$$

Measured soft function contribution (NLO/NLL')

$$f_S(\tau, \mu) = -\frac{\alpha_s(\mu) C_F}{\pi} \frac{1}{1-a} \left\{ \left[\ln \frac{\mu \tan^{1-a} \frac{R}{2}}{\omega \tau} + H(-1-\Omega) \right]^2 + \frac{\pi^2}{6} - \psi^{(1)}(-\Omega) \right\},$$

Apply to Heavy Quarkonium?

Non-relativistic QCD Factorization Formalism

$$\sigma(gg \to J/\psi + X) = \sum_n \sigma(gg \to c\bar{c}(n) + X) \langle \mathcal{O}^{J/\psi}(n) \rangle$$

Expand in α_s

Scaling in v

NRQCD Power Counting in α_s, v		
Mechanism	$d_n(z)$	$\langle \mathcal{O}_n^H \rangle$
${}^3S_1^{(1)}$	α_s^3	v^3
${}^3S_1^{(8)}$	α_s	v^7
${}^1S_0^{(8)}$	α_s^2	v^7
${}^3P_J^{(8)}$	α_s^2	v^7

with $n = {}^{2S+1} L_J^{(1,8)}$

Bodwin, Braaten, Lepage

FJF's and Quarkonia Production

Discriminating power between NRQCD production mechanisms





Baumgart, Mehen, Leibovich, Rothstein, arXiv:1406.2295

FJF's and Quarkonia Production

Discriminating power between NRQCD production mechanisms



Baumgart, Mehen, Leibovich, Rothstein, arXiv:1406.2295

Details of Calculation

Hard, Soft Unmeasured, and Unmeasured Jet Functions

$$H_2(\mu) = 1 - \frac{\alpha_s(\mu) C_F}{2\pi}\left[8 - \frac{7\pi^2}{6} + \ln^2\frac{\mu^2}{\omega^2} + 3\ln\frac{\mu^2}{\omega^2}\right]$$

$$S^{\text{unmeas}}(\mu) = 1 + \frac{\alpha_s(\mu) C_F}{2\pi}\left[\ln^2\frac{\mu^2}{4\Lambda^2} - \ln^2\frac{\mu^2}{4\Lambda^2 r^2} - \frac{\pi^2}{3}\right]$$

$$J_{\bar{n}}^{(\bar{b})}(\mu) = 1 + \frac{\alpha_s(\mu) C_F}{2\pi} J^q_{\text{alg}}(\mu).$$

RG Evolution Factor

$$\Pi(\mu, \mu_H, \mu_\Lambda, \mu_{J_{\bar{n}}}, \mu_{J_n}, \mu_{S^{\text{meas}}}) = \prod_{i=H, J_{\bar{n}}, S^{\text{unmeas}}} \exp(K_i(\mu, \mu_i)) \left(\frac{\mu_i}{m_i}\right)^{\omega_i(\mu,\mu_i)}$$

$$\times \frac{1}{\Gamma(-\Omega(\mu_{J_n}, \mu_{S^{\text{meas}}}))} \times \prod_{i=J_n, S^{\text{meas}}} \exp(K_i(\mu, \mu_i) + \gamma_E \omega_i(\mu, \mu_i)) \left(\frac{\mu_i}{m_i}\right)^{j_i \omega_i(\mu,\mu_i)}$$

Resumming Logarithms

$$
\begin{aligned}
A = a &+ \alpha \left(b_1 + b_2 \log\left(\frac{\mu}{\mu_0}\right)\right) \\
&+ \alpha^2 \left(c_1 + c_2 \log\left(\frac{\mu}{\mu_0}\right) + c_3 \log^2\left(\frac{\mu}{\mu_0}\right)\right) \\
&+ \alpha^3 \left(d_1 + d_2 \log\left(\frac{\mu}{\mu_0}\right) + d_3 \log^2\left(\frac{\mu}{\mu_0}\right) + d_4 \log^3\left(\frac{\mu}{\mu_0}\right)\right) \\
&+ \alpha^4 \left(e_1 + e_2 \log\left(\frac{\mu}{\mu_0}\right) + e_3 \log^2\left(\frac{\mu}{\mu_0}\right) + e_4 \log^3\left(\frac{\mu}{\mu_0}\right) + e_5 \log^4\left(\frac{\mu}{\mu_0}\right)\right) \\
&+ \ldots
\end{aligned}
$$

N³LL **NNLL** **NLL** **LL**

$$N^{n-m}LL \sim \sum \alpha_s^n \log^m\left(\frac{\mu}{\mu_0}\right)$$

Resummation of Logarithms

Evolve each function to common scale using RG



J/ψ Production Mechanisms

Diagrams for each singlet/octet channels





Characteristic Scales in Factorization Theorem



EFT counting	matching/ matrix element	$\Gamma_{\rm cusp}$	$\gamma_{H,J,S}$	$\beta[\alpha_s]$
LL	tree	1-loop	tree	1-loop
NLL	tree	2-loop	1-loop	2-loop
NNLL	1-loop	3-loop	2-loop	3-loop

S.D.Ellis, et. al, arXiv:1001.0014

Global Fits to World's Data

Fit done on 194 data points, 26 data sets



Buttenschon, Kniehl (2011), arXiv:1105.0820

Attempts to Fix Polarization Problem

Simultaneous NLO fit to CMS, ATLAS high p_t production, polarization







Chao, et al. (2012),arXiv:1201.2675

$\langle\mathcal{O}(^3S_1^{[1]})\rangle$ GeV^3	$\langle\mathcal{O}(^1S_0^{[8]})\rangle$ 10^{-2}GeV^3	$\langle\mathcal{O}(^3S_1^{[8]})\rangle$ 10^{-2}GeV^3	$\langle\mathcal{O}(^3P_0^{[8]})\rangle/m_c^2$ 10^{-2}GeV^3
1.16	8.9 ± 0.98	0.30 ± 0.12	0.56 ± 0.21
1.16	0	1.4	2.4
1.16	11	0	0

Inconsistent with global fits!

NRQCD Fragmentation Functions

Matching QCD and NRQCD



Perturbatively Calculable Frag. Functions

$$D^{^3S_1^{(8)}}_{g\to J/\psi}(z,2m_c) = \frac{\pi\alpha_s(2m_c)}{24m_c^3}\langle\mathcal{O}^{J/\psi}(^3S_1^{(8)})\rangle\delta(1-z)$$

Braaten, Chen, hep-ph/9610401

Braaten, Chen, hep-ph/9604237

Braaten, Yuan, hep-ph/9302307

Definitions of Operators

QCD Fragmentation Function

$$D^h_{q,\mathrm{bare}}(x) = \frac{1}{x}\int \mathrm{d}^2 p_h^\perp \int \frac{\mathrm{d}y^+\,\mathrm{d}^2 y_\perp}{2(2\pi)^3}\, e^{ik^- y^+/2} \sum_X \frac{1}{2N_c}\,\mathrm{tr}\Big[\frac{\bar{\not n}}{2}\,\langle 0|\psi(y^+,0,y_\perp)|Xh\rangle\langle Xh|\bar\psi(0)|0\rangle\Big]$$

SCET Fragmentation Function

$$D_q^h(\frac{p_h^-}{\omega},\mu) = \pi\omega \int dp_h^+ \frac{1}{4N_c}\,\mathrm{Tr}\sum_X \bar{\not n}\,\langle 0|\,\delta_{\omega,\bar{\mathcal{P}}}\delta_{0,\mathcal{P}_\perp}\chi_n(0)\,|Xh\rangle\,\langle Xh|\,\bar\chi_n(0)\,|0\rangle$$

SCET Jet Function

$$J(p^\mu) = \frac{1}{8\pi N(\bar n\cdot p)}\sum_X \int d^4x e^{ipx}\,\mathrm{Tr}\left[\langle\Omega|\,\bar\chi_n(x)\,|X_n\rangle\,\langle X_n|\,\bar{\not n}\chi_n(0)\,|\Omega\rangle\right]$$

SCET Fragmenting Jet Function

$$\mathcal{G}^h_{q,\mathrm{bare}}(s,z) = \int \mathrm{d}^4 y\, e^{\mathrm{i}k^+ y^-/2}\int \mathrm{d}p_h^+ \sum_X \frac{1}{4N_c}\,\mathrm{tr}\left[\frac{\bar{\not n}}{2}\langle 0|[\delta_{\omega,\bar{\mathcal{P}}}\,\delta_{0,\mathcal{P}_\perp}\chi_n(y)]|Xh\rangle\langle Xh|\bar\chi_n(0)|0\rangle\right]$$